



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### **Divergence-time estimates for hominins provide insight into encephalization and body mass trends in human evolution**

**Citation for published version:**

Püschel, HP, Bertrand, OC, O'reilly, JE, Bobe, R & Püschel, TA 2021, 'Divergence-time estimates for hominins provide insight into encephalization and body mass trends in human evolution', *Nature Ecology & Evolution*. <https://doi.org/10.1038/s41559-021-01431-1>

**Digital Object Identifier (DOI):**

[10.1038/s41559-021-01431-1](https://doi.org/10.1038/s41559-021-01431-1)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Nature Ecology & Evolution

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



**Divergence-time estimates for hominins provide insight into encephalization and body mass trends in human evolution**

Hans P. Püschel<sup>1\*</sup>, Ornella C. Bertrand<sup>1</sup>, Joseph E. O'Reilly<sup>2</sup>, René Bobe<sup>3,4</sup>, Thomas A. Püschel<sup>3\*</sup>

<sup>1</sup> School of GeoSciences, University of Edinburgh, Grant Institute, James Hutton Road, Edinburgh, EH9 3FE, Scotland, United Kingdom.

<sup>2</sup> MRC Human Genetics Unit, MRC Institute of Genetics & Molecular Medicine, University of Edinburgh, Western General Hospital, Crewe Road, Edinburgh, EH4 2XU, Scotland, United Kingdom.

<sup>3</sup> Primate Models for Behavioural Evolution Lab, Institute of Cognitive and Evolutionary Anthropology, School of Anthropology, University of Oxford, 64 Banbury Road, Oxford, OX2 6PN, England, United Kingdom.

<sup>4</sup> Gorongosa National Park, Sofala, Mozambique.

\*Corresponding authors:

Hans P. Püschel

E-mail: H.P.Puschel-Rouliez@sms.ed.ac.uk

Phone: +44 (0) 7741054298

Thomas A. Püschel

E-mail: thomas.puschelrouliez@anthro.ox.ac.uk

Phone: +44 (0) 7476608464

## Abstract

Quantifying speciation times during human evolution is fundamental as it provides a timescale to test for the correlation between key evolutionary transitions and extrinsic factors like climatic or environmental change. Here we applied a Total Evidence Dating approach to a hominin phylogeny to estimate divergence times under different topological hypotheses. The time scaled phylogenies were subsequently used to perform ancestral state reconstruction of body mass and phylogenetic encephalization quotient (PEQ). Our divergence time estimates are consistent with other recent studies that analysed extant species. We show that the origin of *Homo* most likely occurred between 4.30 and 2.56 Ma. The ancestral state reconstructions show a general trend towards a smaller body mass before the emergence of *Homo*, whilst followed by a trend towards a greater body mass. PEQ estimations display a general trend of gradual but accelerating encephalization evolution. The obtained results provide a rigorous temporal framework for human evolution.

## Introduction

Establishing an evolutionary timescale for human evolution is of essential relevance in palaeoanthropology<sup>1–3</sup> because reliable estimates of the timing of speciation events across the hominin phylogeny facilitate the correlation of these events with both abiotic and biotic processes on geological timescales. An accurate timescale also provides a framework to test for associations between landmark evolutionary changes and different putative extrinsic causal factors such as changes in climate or other environmental influences<sup>4,5</sup>. Despite recent relevant fossil findings, the antiquity and emergence of the genus *Homo*, as well as the timing of the divergence of our lineage with other African apes have not found a consensus<sup>6–9</sup>. Traditionally, palaeoanthropologists have employed maximum parsimony analysis to infer hominin phylogenetic relationships<sup>10–12</sup>, but this approach does not explicitly estimate

divergence-times as part of the estimation of the phylogeny. Previous studies<sup>13,14</sup> computed confidence intervals for the disappearances and appearances of several hominins. However, their approach is not easily applicable to several available hominin taxa, as it requires the availability of extensive palaeontological datasets. Additionally, this method only provides confidence intervals for local first and last appearance data, which do not correspond to the global origination and extinction dates of the taxa under analysis. Other studies have computed these values but their results have been limited by either focusing on a restricted number of hominin taxa<sup>2</sup> or because the assumptions of their applied methods were not met<sup>15</sup>. Bayesian phylogenetic inference methods have also been applied to morphological data<sup>16,17</sup> to provide divergence time estimates. However, these estimates were exclusively based on anatomical data, even though it is now widely known that the fragmentary nature of the fossil record is not enough to compute reliable divergence estimates<sup>18</sup>, and that it is necessary to consider the molecular information available for several hominin and ape taxa in the analyses.

There is currently a consensus that the most feasible way of determining an accurate evolutionary timescale is by using the molecular clock, a prospect that has progressively concretised with the development of Bayesian relaxed clock methods<sup>19–22</sup>. Bayesian divergence-time estimates require the use of prior probability distributions to incorporate fossil evidence for calibrating the tree. Recently, a new approach known as ‘total-evidence dating’ (TED), ‘tip-dating’ or ‘integrative dating’<sup>23</sup> has been developed<sup>24,25</sup>. TED complements the molecular sequence data derived from extant species with morphological information from both living and extinct species, which allows a more thorough inclusion of fossil data in the analysis and estimation of divergence-times.

Hence, in this work we applied this divergence-time estimation method to produce total evidence evolutionary timescales for the hominin clade. We considered four different topological hypotheses with alternative reasonable affinities for problematic hominin taxa (Fig. 1). This is highly relevant because these timescales can be used for dating the origin of *Homo* or any other hominin genus, inferring evolutionary rates and patterns, as well as providing a better understanding of the coevolution of hominins and their environment. Additionally, we subsequently used the dated trees to carry out ancestral state reconstruction (ACSR) of two evolutionary important phenotypic characters: body mass and phylogenetic encephalization quotient (PEQ). Body mass impacts almost every aspect of an animal's biology and ecology<sup>26,27</sup>, hence its importance in any palaeobiological inference, whilst an evolutionary trend of increasing encephalization is one of the hallmark processes in human evolution<sup>28,29</sup>.

## Results

The TED analyses (Fig. 2, Extended Data Fig. 1 and Table 1) show, in general, better-resolved divergence times for the nodes that have both morphological and molecular information available, in comparison with the nodes estimated with morphological information alone. The common ancestor of hominins and *P. troglodytes* (node 2) was dated at around 7.5 Ma with an uncertainty range of 8.59 – 6.61 Ma considering the 95% highest posterior density intervals (HPD) of all the trees. The common ancestor of the genus *Homo* (node 13 or 14 for *H. floresiensis* hypothesis) was dated at around 3.3 Ma with an uncertainty range of 4.30 – 2.56 Ma considering the 95% HPD of all the trees. In general, estimations for this node were consistent between the four trees, although the biggest difference resulted when *H. floresiensis* is removed from the base of *Homo*, making the node's age slightly younger (Fig. 2d). The prior sensitivity analysis (Extended Data Fig. 2) shows mostly minor

differences between divergence-times in the original analysis and analyses using different priors in relevant parameters, suggesting that the TED analyses are robust to changes in the employed priors.

The maximum likelihood (ML) ACSR based on the four consensus trees (Figs. 3, 4, Extended Data Figs. 3-6) are consistent with the ACSR based on samples of the posterior trees from these analyses (Fig. 5 and Extended Data Fig. 7). The main difference is an apparent PEQ overestimation in the case of the ACSR based on the consensus trees versus the ACSR based on samples of the posterior trees (Fig. 5). These differences are explained because using equations based on PGLS regressions of the consensus trees (Extended Data Fig. 8) returned lower expected brain masses calculated for the trees' tips on average than equations based on the sampled trees (Extended Data Fig. 9). However, the general PEQ trends remain the same. Similarly, the brain mass ACSR versus body mass ACSR regressions results show a consistently similar pattern for the four analyses. When we analyse all the nodes together in the four trees, the slopes are slightly positive and the  $R^2$  values are low (around 0.11) (Extended Data Fig. 10a-d). However, when we split the data there are clearly two different trends. Before node 13/14 the slopes are negative with a moderate  $R^2$  (around 0.46) (Extended Data Fig. 10e-h), whereas after node 13/14 the slopes are strongly positive with a high  $R^2$  (around 0.96) (Extended Data Fig. 10i-l).

The speciation events that occurred since the divergence from *G. gorilla* to the common ancestor of *Homo* (nodes 1-9 and 13/14), all occurred within the latest Miocene and Pliocene (Fig. 2 and Table 1). For that period, the body mass ACSR show that from a common ancestor of around 71 kg at 10.4 Ma, there is a trend of rapid decrease in body mass reaching around 38 kg by 3.3 Ma (Figs. 3, 5a-d, and Extended Data Fig. 5). That is 1.87 smaller in 7.1 Myr. However, the PEQ ACSR show an opposite trend, increasing from around 0.87 to 1.88

in the same period (Figs. 4, 5e-h and Extended Data Fig. 6), which is 2.16 times greater. The effect of removing of *H. floresiensis* from the base of *Homo* (Figs. 4d, 5h and Extended Data Fig. 6d) is a higher estimated PEQ for the last common ancestor of this genus (i.e., 2.16) and the surrounding nodes. Following the path leading to *Au. africanus* and the members of the genus *Paranthropus* (nodes 1-10) the body mass tends to decrease to around 37 kg at node 10, which means a reduction of 1.92 times from the root in 6.6 Myr. Nevertheless, the path from nodes 9 to 10 does not indicate an increase in PEQ. Instead, it corresponds to the beginning of a slight reduction in PEQ in the lineage leading to *Paranthropus*. Interestingly, the last member of this genus, *P. robustus*, displays an increase in PEQ just before the extinction of the lineage. The incorporation of *Au. sediba* as the sister taxon of *Au. africanus* did not have a substantial influence on the ACSR of the neighbouring nodes.

Speciation events since the common ancestor of *Homo* start to occur around the mid-Pliocene at 3.3 Ma and end around the mid-Pleistocene at 0.6 Ma (Fig. 2 and Table 1). In contrast to the previous decreasing trend in body mass, the body mass ACSR show an increment from around 38 kg to 65 kg in this period (nodes 13/14-24; Figs. 3, 5a-d, and Extended Data Fig. 5). This corresponds to an increase in body mass of 1.71 times in 2.7 Myr, which means a reversion in body size to similar levels as the one observed at the tree's root but in less than half of the time. In other words, the rate of body mass evolution approximately doubled in the genus *Homo* from what has been previously seen (10.0 kg/Myr vs 4.6 kg/Myr). However, these increments were not equal in all nodes and some groups even show a body mass reduction. For instance, the parallel branch that goes from node 14 to *Homo habilis* shows a decrease in body mass regardless of the position of *Au. sediba*. The effect of removing *Au. sediba* from *Homo* results in a slight increase in the reconstructed body mass at node 14. Similarly, *H. floresiensis* and *H. naledi* also displayed a considerable decrease in body mass from their ancestors in all the hypotheses. However, the decrease and its rate vary with the

different hypotheses. When *H. floresiensis* is positioned at the base of *Homo*, the decrease is around 10 kg (-3.0 kg/Myr), while when it is considered the sister taxon of the Asian *H. erectus* the decrease is approximately 22 kg (-13.0 kg/Myr). When *H. naledi* is at the stem of *H. antecessor* the decrease is around 13 kg (-10.1 kg/Myr), whereas when it is the sister taxon of the African *H. erectus* the decrease is approximately 5 kg (-3.0 kg/Myr).

The PEQ ACSR for the period since the common ancestor of *Homo* (node 13/14) keeps the previously observed increasing trend, augmenting 1.53 times in 2.70 Myr, from around 1.88 in node 13/14 to 2.88 in node 24 (Figs. 4, 5e-h and Extended Data Fig. 6). Even though the increase in PEQ is similar with the PEQ evolution before the common ancestor of *Homo* (1.00 and 1.01 respectively), its rate increased 2.64 times, going from 0.14 PEQ/Myr to 0.37 PEQ/Myr. Within this general PEQ evolutionary trend, there are three taxa that stand out from the rest: *H. floresiensis*, *H. naledi* and *H. sapiens*, although in the first two species the specific trends vary with the different hypotheses. When *H. floresiensis* is positioned is at the base of *Homo*, it displays a slight decrease or stasis in PEQ from its ancestor exhibiting a value around 1.7 for 3.4 Myr of evolutionary history, while when it is considered the sister taxon of the Asian *H. erectus* the decrease is 0.52 in 1.7 Myr (-0.30 PEQ/Myr). When *Homo naledi* is at the stem of *H. antecessor*, the PEQ decreases around 0.85 in 1.3 Myr (-0.65 PEQ/Myr), whereas when it is the sister taxon of the African *H. erectus* the decrease is approximately of 0.69 in 1.7 Myr (-0.42 PEQ/Myr). This strong decrease under different scenarios puts *H. naledi* very close to the PEQ values for *Au. afarensis*. In contrast, *H. sapiens* stands out due to its rapid increase in PEQ from its common ancestor with *H. neanderthalensis* (node 24), going from around 2.88 to 3.22 in approximately 0.55 Myr. That is 0.62 PEQ/Myr, which is 1.68 times greater compared to the rate observed since the common ancestor of *Homo* (nodes 13/14 to 24) between 3.3 Ma and 0.6 Ma, and 4.43 greater



in comparison to the rate observed since the common ancestor of *G. gorilla* and the ancestor of *Homo* (nodes 1 to 13/14) between 10.4 Ma and 3.3 Ma.

## Discussion

We have presented here TED estimates of the divergence-times of most hominin taxa under different hypotheses. Our divergence estimates are in general agreement with previous molecular studies using fossil node-calibrations<sup>30–34</sup>, as well as with fossil calibration independent methods, such as those using generation times<sup>35</sup>. The topology of our trees (Fig. 2) differs from the trees of Dembo *et al*<sup>16,17</sup> in the position of *H. neanderthalensis* relative to *H. sapiens* and *H. heidelbergensis*, which is likely due to the influence of the mtDNA in our Total Evidence analysis. The position of these groups plus Denisovans is the same as obtained in a previous study<sup>34</sup> using mtDNA but differs from studies using nuclear DNA<sup>36</sup>, probably related to a mtDNA introgression event that occurred ~270 ka<sup>37</sup>. We think our divergence-time estimates are robust and consistent with the current evidence available. However, further refinements in some of the fossilized birth-death (FBD) model's assumptions, namely being able to consider a nonuniform fossil sampling among clades<sup>38</sup>, and new fossil discoveries, could further improve these estimates.

Our ACSR results from the four different considered hypotheses are consistent between them, showing that refining the phylogenetic affinities of problematic taxa (e.g., *H. naledi*) would probably have a minor impact in major body size and encephalization trends in hominin evolution. Furthermore, our ACSR results are consistent with previous studies that show that body size in hominins has not been a simple linear increment since the divergence with *P. troglodytes*<sup>26,27,39</sup>. The observed decreasing trend in body size from the root of the tree is in agreement with previous studies that suggest a chimpanzee-sized common ancestor

with *P. troglodytes*<sup>40,41</sup>. Our ACSR results also displayed a general trend towards greater body size that started after the emergence of the last common ancestor of *Homo*, contrary to a previous claim stating that there were no clear body mass temporal trends in hominin evolution<sup>26</sup>. Instead, our results showed a complex history of body size changes that do not correlate linearly with brain size changes. It may be that after the emergence of the genus *Homo*, brain size carried body size increases as previously suggested<sup>42</sup> (Extended Data Fig. 10i-l). However, before the emergence of *Homo*, brain size and body size seem to be decoupled or following opposite directions (Extended Data Fig. 10e-h). Our results also agree with Pagel's<sup>43</sup> seminal work and other studies (e.g.,<sup>44-46</sup>) that showed a general trend of gradual but accelerating brain size evolution in hominins. However, some of them (i.e.,<sup>44,45</sup>) directly analysed endocranial volume without taking into account body mass estimations, which we considered in our PEQ estimations. This is problematic because it is only when body size is considered<sup>47</sup> and phylogeny is included that we can consider whether observed brain size differences are significant in terms of encephalization<sup>28,48,49</sup>. We are confident that our ACRS are reliable and consistent with the current evidence. Nevertheless, it is important to note that the inclusion of more fossils near the root could have an impact in the ACSR, in particular for the oldest nodes of the tree. Additionally, ACSR could be affected by the use of different methods and models of character evolution<sup>50</sup>.

The fact that PEQ evolution tends to accelerate when moving from a relatively stable climate to more unstable climatic conditions (Fig. 2) appears to agree with the 'variability selection' hypothesis which correlates major adaptations in hominins with periods of high climatic variability<sup>51,52</sup>. Before the emergence of *Homo*, hominin evolution occurred in a mostly warm period although with a global cooling trend, which had started after the Mid-Miocene Climatic Optimum and the recovery of Antarctic ice-sheets around 10 Ma<sup>53</sup>. From a common ancestor with *P. troglodytes* at around 7.5 Ma, bipedalism started to emerge at least by 6

214 Ma<sup>54–57</sup> in the hominin lineage. This relatively stable period was interrupted by considerable  
215 temperature oscillations around the Miocene/Pliocene boundary at 5.3 Ma. Subsequently with  
216 the development of 23 kyr dominant glacial cycles, we observed the emergence of the genus  
217 *Homo* and an evolutionary shift, displaying the end of a general trend towards smaller body  
218 size and the beginning of an acceleration of the increase in PEQ and the start of a general  
219 trend towards larger body masses (Figs. 2-4). Even though the earliest member of *Homo*  
220 discovered so far was from 2.8 Ma<sup>58</sup>, our analysis allows us to predict the presence of early  
221 representatives of the *Homo* lineage not yet found (or identified) in the African fossil record  
222 around 3.3 Ma (i.e., 0.5 Myr earlier). Interestingly, 3.3 Ma is the age attributed to stone tools  
223 discovered in West Turkana<sup>59</sup>, which are commonly associated with *K. platyops*<sup>60</sup> and *Au.*  
224 *afarensis*<sup>61</sup>.

225 After the Pliocene-Pleistocene boundary at 2.58 Ma, the trend towards cooler temperatures  
226 and aridity continues, and 41 kyr dominant glacial cycles are established, with an  
227 intensification of climatic fluctuations. As PEQ continues to increase, so does the evidence of  
228 increasingly complex behaviours: tool innovations (Oldowan [2.6 Ma<sup>62</sup>], Acheulean [1.7-1.4  
229 Ma<sup>63</sup>] and Aurignacian at [120–~50 ka<sup>64</sup>], the use of fire (from 1.5 Ma onwards<sup>65,66</sup> [strong  
230 evidence at 1.0–0.5 Ma<sup>67</sup>]), cooking and more frequent meat consumption<sup>68–70</sup> and the ability  
231 to produce art at ~540–430 ka<sup>71</sup>. By 300 ka Africa was inhabited by at least three *Homo*  
232 species, *H. sapiens*, *H. heidelbergensis* and *H. naledi*, and Eurasia by *H. neanderthalensis*,  
233 Denisovans, *H. floresiensis*, *H. luzonensis* and also possibly *H. erectus* and *H.*  
234 *heidelbergensis*<sup>72</sup>. *H. sapiens* is the hominin species with the highest PEQ, so a possible  
235 explanation for its exclusive continuation would be that this difference in PEQ allowed *H.*  
236 *sapiens* to outcompete its contemporary relatives<sup>47,73–75</sup>. Even though it has been recently  
237 established that there was not a unique ‘Out of Africa’ event in *H. sapiens* history<sup>76–79</sup>, it is  
238 widely accepted that Neanderthals were eventually displaced by *H. sapiens* in Europe by ~39

239 ka<sup>80</sup>. However, it is now understood that *H. neanderthalensis* was capable of very complex  
240 human behaviours<sup>81,82</sup>.

241 Like *H. sapiens*, *H. neanderthalensis* was certainly a cultural niche constructor<sup>83–85</sup> under  
242 harsh glacial-interglacial temperature fluctuations<sup>86</sup>. Nonetheless, *H. sapiens* not only  
243 displays a higher PEQ, but also a higher rate of change in PEQ compared to *H.*  
244 *neanderthalensis*. This probably means that within Hominini, PEQ selection was particularly  
245 strong within and between the metapopulations of *H. sapiens*<sup>87</sup> in an arid-moist fluctuating  
246 Africa<sup>86</sup>. A larger relative brain mass has been associated in mammals with behavioural  
247 flexibility, adaptation and resilience in variable environmental conditions<sup>88</sup>. Therefore in spite  
248 of the fact that the behavioural gap between the two species may have been minimal, even a  
249 small advantage in terms of behavioural flexibility and ability to adjust in a variable  
250 environment as it was during the late Pleistocene<sup>89</sup>, could have had enormous benefits in  
251 terms of fitness and successful competition for *H. sapiens*<sup>90,91</sup>. A similar explanation could  
252 also be applied for the demise of the rest of our contemporaneous relatives exhibiting an even  
253 lower PEQ.

254 Considering the evolution of *H. floresiensis*, it seems that selection was acting on body size  
255 by means of heterochrony<sup>92,93</sup> favouring a reduction in body size as a mean of decreasing the  
256 energetic expenditure in a small island environment with limited resources as it also probably  
257 occurred for *Stegodon florensis*<sup>94–96</sup>. However, our results show that resolving the  
258 phylogenetic affinities of *H. floresiensis* would have important implications for the  
259 evolutionary trends on this taxon. The body size reduction in *H. floresiensis* is more  
260 spectacular if this taxon derives or is considered closely related to Asian *H. erectus*<sup>97</sup>, a  
261 scenario that also implies a notorious encephalization reduction. Nevertheless, more recent  
262 studies favour the position of *H. floresiensis* at the base of *Homo*<sup>16,17,98</sup>, which would then

favour a encephalization stasis scenario. Interestingly, if tool development can be associated with a certain level of cognition, considering the tools attributed to *H. floresiensis*<sup>99,100</sup>, we anticipate that the common ancestor of *Homo* with a similar PEQ value, was most likely able make stone tools as well.

Previous studies have described *H. naledi* as a small bodied and small brained hominin of the genus *Homo*<sup>101</sup>, with cranial<sup>102</sup>, endocranial<sup>103</sup> and postcranial features<sup>104,105</sup> that support this placement. There are two hypotheses which attempt to explain this small brain as either a (1) retention from the common ancestor from the genus *Homo*, or (2) a reduction from a later big-brained form of *Homo*<sup>103</sup>. The PEQ trends displayed by *H. naledi*'s ACSR supports the second hypothesis because there is an extraordinary reduction in PEQ from a big-brained ancestor in a relatively short time, although this reduction is faster if *H. naledi* is at the stem of *H. antecessor* as previously suggested<sup>17</sup>. *H. naledi* lived between 236 ka and 335 ka<sup>106</sup> in South Africa, with a PEQ around 1.5, which is really close to *Au. afarensis* and other australopiths. This happened at a time in which big bodied and big brained hominins were the norm in continental landmasses, like *H. sapiens* and *H. neanderthalensis* both with a PEQ over 2.7. Even the insular small sized *H. floresiensis* had a higher PEQ. In a context of increasing PEQ over the hominin lineage at that time, the PEQ reduction in *H. naledi* could perhaps be explained, by the specialization in a niche of scarce and/or low energy food resources in which an expensive large brain would be prejudicial. Hypotheses in relation to the cost of encephalization like the 'expensive tissue'<sup>107,108</sup> and the 'energy trade off' hypotheses<sup>109</sup>, could potentially explain this trend in *H. naledi*. Less energy expenditure could also explain the considerable body size reduction observed in *H. naledi* from its reconstructed bigger-brained ancestors. In fact, through dental topography comparisons it has been suggested that *H. naledi* was occupying a distinct ecological niche, which was different from previous and contemporaneous hominins<sup>110</sup>.

In conclusion, our TED analyses and ACSR results showed that (1) the last common ancestor of the genus *Homo* most likely appeared around 3.3 Ma (between 4.30 and 2.56 Ma) with a body size close to that of *Au. afarensis* and an encephalization very similar to *H. floresiensis*, (2) hominin body mass evolution followed a general decreasing trend before the emergence of *Homo* and exhibited a general increasing trend afterwards, (3) hominins displayed a general trend of gradual but accelerating encephalization through time.

## Methods

### Data collection and Total Evidence analyses

We used TED analyses which are a collection of Bayesian phylogenetic methods (see<sup>111</sup> for a general primer and <sup>19,20,22,112,113</sup> for reviews of Bayesian molecular dating methodology). For the TED analysis, our taxon sampling was similar to previous published analyses<sup>17</sup> but Denisovans were also included. The morphological data were obtained from the same study<sup>17</sup>, and comprised a supermatrix of 391 craniodental characters from matrices used in 13 previous studies<sup>114–126</sup>. Even though there are more recently published hominin phylogenetic analyses computed using different morphological matrices<sup>127,128</sup>, the morphological matrix used here<sup>17</sup> is the most complete, to our knowledge, in terms of the number character states and hominin species included.

The molecular data were complete mtDNA genomes extracted from GenBank for the species for which it was available: *Gorilla gorilla* (KF914214.1), *Pan troglodytes* (JF727180.2), *Homo heidelbergensis* (KF683087.1), *Homo neanderthalensis* (MK123269.1), *Homo sapiens* (KC417443.1) and Denisovans (KX663333.1). Following previous analyses<sup>34,129</sup>, we removed the D-loop region from the mtDNA due to the differential rate at which it acquires substitutions. We aligned the sequences with the MUSCLE algorithm in MEGA X<sup>130</sup>. Then, we analysed the alignment using PartitionFinder2<sup>131</sup> using the “greedy” algorithm<sup>132</sup>, in order

312 to select the most appropriate models of molecular evolution for the different protein coding  
313 and non-coding regions of the mtDNA. The best partitioning scheme on the basis of AICc  
314 score included 16 partitions for the mtDNA (Supplementary Table 1). We used these  
315 partitions for the mtDNA sequences, and the Mkv+  $\Gamma$  model<sup>133</sup> for the morphological data,  
316 unlinking the model parameters across these partitions. In order to avoid a mismatch in our  
317 model, we used the dates associated with our data following the recommended procedure<sup>134</sup>,  
318 which meant that we did not necessarily calibrate a tip using the oldest or first fossil  
319 occurrence for a particular taxon. Hence, we calibrated the fossil tips of the tree using the age  
320 of the fossil specimen used for coding morphology in taxa without mtDNA available. In taxa  
321 with mtDNA sequences available, the sequences were selected from individuals aged equally,  
322 or as close as possible, to the morphologically coded fossils, and the age associated with  
323 these sequences was used to calibrate the fossil tips. We also considered radiometric age  
324 uncertainties using a uniform distribution between the maximum and minimum estimated  
325 ages for each fossil when available<sup>112</sup> (further information can be found in Supplementary  
326 Table 2).

327 The selected outgroup was *Gorilla gorilla* and the root of the tree was calibrated using a  
328 uniform distribution between 10.00 and 12.5 Ma. The minimum age of 10 Ma and the  
329 maximum age of 12.5 were based on the appearance of the proposed stem member of the  
330 gorilla clade *Nakalipithecus nakayamai*<sup>135,136</sup> and the probable crown pongine  
331 *Sivapithecus*<sup>137,138</sup>, respectively. Even though there is uncertainty regarding the exact  
332 phylogenetic placement of *N. nakayamai*<sup>135,136</sup>, we consider that the anatomical features  
333 linking it with gorillas are strong enough to use the age associated with this taxon as  
334 minimum divergence date for hominines, particularly when considering the possible  
335 ancestral-descendent relationship between *N. nakayamai* and the basal gorillin  
336 *Chororapithecus abyssinicus*<sup>135,139,140</sup>.

We used a normally distributed clock rate prior, with a mean of 0.025 and standard deviation of 0.05, which is consistent with previous estimates of the mitochondrial rate of evolution in humans<sup>141,142</sup>. The Independent Gamma Rate (IGR) relaxed clock model was used for modelling branch rate variation, employing a clock rate variance prior with an exponential distribution of rate 10. We used the FBD model as the prior on divergence times, using an exponential net diversification prior with rate 1, a beta turnover prior with shape parameters  $\alpha=1$  and  $\beta=1$ , a beta fossil sampling proportion prior with shape parameters  $\alpha=1$  and  $\beta=1$  and an extant sampling proportion of 1. The priors employed in clock rate variance and the FBD model were intentionally diffuse, reflecting the general uncertainty in our prior expectation of the distribution of these parameters.

Even though our main focus was estimating hominin divergence-times, we are aware that that there is still controversy regarding the phylogenetic placement of some of the taxa included in our matrix, particularly in the placement of *Australopithecus sediba*<sup>1,117</sup>, *Homo floresiensis*<sup>16,97,98</sup> and *Homo naledi*<sup>17,143</sup>. Therefore, we considered four different topological hypotheses for constraining four independent phylogenetic analyses (Fig. 1): a), a topology similar to the phylogeny of the analysis from which we extracted the morphological data<sup>17</sup>; b), the same topology as a) but moving *Au. sediba* from the *Homo* clade, to be the sister taxa of *Australopithecus africanus* as has been recently suggested<sup>144</sup>; c), the same topology as a) but changing the position of *Homo naledi* from the stem of *Homo antecessor* to be the sister taxa of the African *Homo erectus* as previously suggested<sup>143,145</sup>; d), the same topology of a) but taking *H. floresiensis* from the base of the genus *Homo* to be the sister taxa of the Asian *Homo erectus* as was originally suggested<sup>97</sup>. Nevertheless, we left *H. sapiens*, *H. neanderthalensis*, *H. heidelbergensis* and Denisovans unconstrained in the four analyses as they had mtDNA sequences which could also be informative of their phylogenetic relationships. Therefore, the constraints we used in all the analyses were soft, so the latter



taxa could be accommodated in any position of the tree according to their morphological, molecular and stratigraphic information.

We performed the analyses with MrBayes 3.2.7a<sup>146</sup> using two runs of four chains and 60 million MCMC generations with the first 25% of samples discarded as burn-in. The analyses were run on CIPRES portal v3.3<sup>147</sup>. We ensured that the average standard deviation of split frequencies was below 0.01 and that all parameters had an effective sample size of more than 200. Additionally, we visually inspected that the two independent runs achieved convergence and stationarity using the program Tracer v1.7.1<sup>148</sup>.

### **Evaluating the prior sensitivity of divergence-time estimates**

In order to evaluate the effect of alternative priors on our divergence-time estimates, we conducted sensitivity analyses in the first tree (i.e., Dembo *et al.*<sup>17</sup> hypothesis) in which we changed the prior distribution of one important model parameter at a time into a reasonable alternative prior distribution, keeping the rest of the model parameters unchanged. This was done for the clock rate variance, the net diversification prior of the FBD model and root age prior. We did this in the former two cases because we wanted to test the effect of using more constrained priors, and in the latter, to account for the possibility that *N. nakayamai* could be a stem hominid before the gorilla–human split<sup>135,136</sup>. Therefore, for the clock rate variance prior, we ran a strict clock and a non-clock analysis constrained to have the same topology. Then, following Ronquist's *et al.*<sup>25</sup> methodology to estimate rate variance, the clock rate variance prior was estimated as 25.04. For the net diversification prior we followed Zhang *et al.*<sup>149</sup> in using an exponential distribution of rate 100. In the case of the root age prior we used a uniform distribution changing the minimum to 8 but keeping the maximum in 12.5 Ma, because of the 8 Ma estimated for appearance of the proposed gorillin *C. abyssinicus*<sup>135,139</sup>,

which have been used to date the minimum age of the gorilla–human split in previous studies<sup>32,33</sup>.

### **Estimating PEQ using a PGLS regression**

The encephalization quotient (EQ) is commonly used to determine how brain size scales with respect to body size for a given individual<sup>48,150–152</sup>. However, EQ does not take into account phylogenetic information, so a newly proposed measurement termed PEQ has been advanced as a way of considering the phylogenetic non-independence between data points<sup>28</sup>.

Body mass and endocranial volume (ECV) were obtained from the literature (see Supplementary Table 3 for further information). When more than one specimen was available, arithmetic averages for body mass and endocranial volume were used. Endocranial volume was converted into brain mass by dividing ECV by 1.036<sup>153</sup>. We used R version 3.6.1<sup>154</sup> and the packages ‘ape’<sup>155</sup> and ‘nlme’<sup>156</sup> to compute phylogenetic correlations and to fit linear models, respectively. We log-scaled the data, and by assuming a Brownian motion model of evolution we calculated a correlation using the corBrownian() function to then fit a linear model (brain mass was the dependent variable, whilst body mass was the independent one) independently for each one of the four consensus trees obtained in the TED analysis (Extended Data Fig. 8).  $R^2$  were calculated using the R package rr2 v1.0.2 and the function R2.pred()<sup>157</sup>. The resulting equations were used to calculate the expected brain mass (E), and PEQ was calculated as the ratio between the actual estimated brain mass (A) and E ( $PEQ = A/E$ ) for each one of the living and fossil taxa (Supplementary Table 3).

### **PEQ and body mass ancestral character state reconstructions**

We used a ML approach to perform the ACSR at the internal nodes of the four consensus phylogenetic trees for 1) body mass and 2) PEQ (and brain mass) under a Brownian motion

model. This procedure was performed using the fastAnc() and the contMap() functions from the package ‘phytools’ v 0.6-99<sup>158</sup>. *Kenyanthropus platyops* and *Australopithecus garhi* tips were dropped from the ACSR for body mass and PEQ as there were no body mass estimations for these taxa available due to their fragmentary fossil record. Similarly, Denisovans were removed from all ACSR analyses as there are no estimates available for their brain and body mass.

#### **Measuring uncertainty in the ML ancestral character state reconstructions**

In order to measure the uncertainty in our ML ACSR we sampled every 10<sup>th</sup> time-calibrated tree from the posterior after discarding the first 25% as burn-in, which it meant a total of 9002 time-calibrated posterior trees sampled for each one of the four analyses. Then we ran ML ACSR analyses in all of these trees and their internal nodes for 1) body mass and 2) PEQ (and brain mass), using the same methods and R packages described in the previous section. For the PEQ ACSR, we previously ran the PGLS analysis for each one of the sampled trees, so the PEQ values in the trees’ tips were independently estimated for each tree. This allowed us to incorporate uncertainty in our ACSR, and to analyse if the patterns observed in the ACSR for the consensus trees hold or not when we looked at different trees recovered from the posterior.

#### **Measuring the relationship between ACSR of brain mass and body mass**

As brain size increase has been proposed as a driver of body size increase<sup>42</sup>, we carried out regressions between our ACSR of brain mass versus body mass in the four consensus trees to test if there was a pattern that could be consistent with that hypothesis. It is important to consider that we did not directly assess the hypothesis a brain size increase drove body mass evolution but rather evaluate if there was a general pattern that could provide further

information about this issue. Both ACSR datasets were log-scaled before performing the regressions.

#### **Data availability**

All data analysed in this study are available in the Supplementary information (Supplementary Tables 2 and 3) and in a permanent Zenodo (zenodo.org) repository at <https://dx.doi.org/10.5281/zenodo.4537445><sup>159</sup>. Additionally, the data is available in an open access repository at <https://github.com/HansPueschel/Hominin-div-time-evolution><sup>159</sup>.

#### **Code availability**

The code and input files are available in a permanent Zenodo (zenodo.org) repository at <https://dx.doi.org/10.5281/zenodo.4537445><sup>159</sup>. In addition, the code and input files are available in as an open access repository at <https://github.com/HansPueschel/Hominin-div-time-evolution><sup>159</sup>.

#### **References**

1. Du, A. & Alemseged, Z. Temporal evidence shows *Australopithecus sediba* is unlikely to be the ancestor of Homo. *Sci. Adv.* **5**, eaav9038 (2019).
2. Du, A., Rowan, J., Wang, S. C., Wood, B. A. & Alemseged, Z. Statistical estimates of hominin origination and extinction dates: A case study examining the *Australopithecus anamensis–afarensis* lineage. *J. Hum. Evol.* **138**, (2020).
3. Finarelli, J. A. & Clyde, W. C. Reassessing hominoid phylogeny: evaluating congruence in the morphological and temporal data. *Paleobiology* **30**, 614–651 (2004).
4. DeMenocal, P. B. African climate change and faunal evolution during the Pliocene-Pleistocene. *Earth Planet. Sci. Lett.* **220**, 3–24 (2004).

- 454 5. DeMenocal, P. B. Climate and human evolution. *Science* **331**, 540–542 (2011).
- 455 6. Antón, S. C. & Josh Snodgrass, J. Origins and evolution of genus *Homo*: New  
456 perspectives. *Curr. Anthropol.* **53**, S479–S496 (2012).
- 457 7. Harcourt-Smith, W. E. H. Early hominin diversity and the emergence of the genus  
458 *Homo*. *J. Anthropol. Sci.* **94**, 19–27 (2016).
- 459 8. Villmoare, B. Early *Homo* and the role of the genus in paleoanthropology. *Am. J. Phys.*  
460 *Anthropol.* **165**, 72–89 (2018).
- 461 9. Ruiters, D. J. de, Churchill, S. E., Hawks, J. & Berger, L. R. Late Australopiths and the  
462 Emergence of *Homo*. *Annu. Rev. Anthropol.* **46**, 99–115 (2017).
- 463 10. Skelton, R. R. & McHenry, H. M. Evolutionary relationships among early hominids. *J.*  
464 *Hum. Evol.* **23**, 309–349 (1992).
- 465 11. Strait, D. S., Grine, F. E. & Moniz, M. A. A reappraisal of early hominid phylogeny. *J.*  
466 *Hum. Evol.* **32**, 17–82 (1997).
- 467 12. Smith, H. F. & Grine, F. E. Cladistic analysis of early *Homo* crania from Swartkrans  
468 and Sterkfontein, South Africa. *J. Hum. Evol.* **54**, 684–704 (2008).
- 469 13. Bobe, R. & Leakey, M. G. Ecology of Plio-Pleistocene Mammals in the Omo—  
470 Turkana Basin and the Emergence of *Homo*. in *The first humans: origins of the genus*  
471 *Homo* (eds. Grine, F. E., Fleagle, J. G. & Leakey, R. E.) 173–184 (Dordrecht:  
472 Springer, 2009). doi:10.1007/978-1-4020-9980-9\_15.
- 473 14. Bobe, R. & Carvalho, S. Hominin diversity and high environmental variability in the  
474 Okote Member, Koobi Fora Formation, Kenya. *J. Hum. Evol.* **126**, 91–105 (2019).

- 475 15. Maxwell, S. J. The quality of the early hominin fossil record: Implications for  
476 evolutionary analyses. PhD, University College London - Birkbeck, London. (2018).
- 477 16. Dembo, M., Matzke, N. J., Mooers, A. & Collard, M. Bayesian analysis of a  
478 morphological supermatrix sheds light on controversial fossil hominin relationships.  
479 *Proc. R. Soc. B Biol. Sci.* **282**, 20150943 (2015).
- 480 17. Dembo, M. *et al.* The evolutionary relationships and age of *Homo naledi*: An  
481 assessment using dated Bayesian phylogenetic methods. *J. Hum. Evol.* **97**, 17–26  
482 (2016).
- 483 18. Donoghue, P. C. J. & Benton, M. J. Rocks and clocks: calibrating the Tree of Life  
484 using fossils and molecules. *Trends Ecol. Evol.* **22**, 424–431 (2007).
- 485 19. Donoghue, P. C. J. & Yang, Z. The evolution of methods for establishing evolutionary  
486 timescales. *Philos. Trans. R. Soc. B Biol. Sci.* **371**, 20160020 (2016).
- 487 20. Dos Reis, M., Donoghue, P. C. J. & Yang, Z. Bayesian molecular clock dating of  
488 species divergences in the genomics era. *Nat. Rev. Genet.* **17**, 71–80 (2016).
- 489 21. Kumar, S. & Blair Hedges, S. Advances in time estimation methods for molecular  
490 data. *Mol. Biol. Evol.* **33**, 863–869 (2016).
- 491 22. Bromham, L. *et al.* Bayesian molecular dating: opening up the black box. *Biol. Rev.*  
492 **93**, 1165–1191 (2017).
- 493 23. Ronquist, F., Lartillot, N. & Phillips, M. J. Closing the gap between rocks and clocks  
494 using total-evidence dating. *Philos. Trans. R. Soc. B Biol. Sci.* **371**, 20150136 (2016).
- 495 24. Pyron, R. A. Divergence time estimation using fossils as terminal taxa and the origins

496 of lissamphibia. *Syst. Biol.* **60**, 466–481 (2011).

497 25. Ronquist, F. *et al.* A total-evidence approach to dating with fossils, applied to the early  
498 radiation of the hymenoptera. *Syst. Biol.* **61**, 973–999 (2012).

499 26. Grabowski, M., Hatala, K. G., Jungers, W. L. & Richmond, B. G. Body mass estimates  
500 of hominin fossils and the evolution of human body size. *J. Hum. Evol.* **85**, 75–93  
501 (2015).

502 27. Jungers, W. L., Grabowski, M., Hatala, K. G. & Richmond, B. G. The evolution of  
503 body size and shape in the human career. *Philos. Trans. R. Soc. B Biol. Sci.* **371**,  
504 (2016).

505 28. Ni, X., Flynn, J. J., Wyss, A. R. & Zhang, C. Cranial endocast of a stem platyrrhine  
506 primate and ancestral brain conditions in anthropoids. *Sci. Adv.* **5**, (2019).

507 29. Melchionna, M. & Mondanaro, A. Macroevolutionary trends of brain mass in  
508 Primates. *Biol. J. Linn. Soc.* **129**, 14–25 (2020).

509 30. Chatterjee, H. J., Ho, S. Y., Barnes, I. & Groves, C. Estimating the phylogeny and  
510 divergence times of primates using a supermatrix approach. *BMC Evol. Biol.* **9**, (2009).

511 31. Dos Reis, M. *et al.* Phylogenomic datasets provide both precision and accuracy in  
512 estimating the timescale of placental mammal phylogeny. *Proc. R. Soc. B Biol. Sci.*  
513 **279**, 3491–3500 (2012).

514 32. Dos Reis, M. *et al.* Using phylogenomic data to explore the effects of relaxed clocks  
515 and calibration strategies on divergence time estimation: Primates as a test case. *Syst.*  
516 *Biol.* **67**, 594–615 (2018).

- 517 33. Pozzi, L. *et al.* Primate phylogenetic relationships and divergence dates inferred from  
518 complete mitochondrial genomes. *Mol. Phylogenet. Evol.* **75**, 165–183 (2014).
- 519 34. Meyer, M. *et al.* A mitochondrial genome sequence of a hominin from Sima de los  
520 Huesos. *Nature* **505**, 403–406 (2014).
- 521 35. Langergraber, K. E. *et al.* Generation times in wild chimpanzees and gorillas suggest  
522 earlier divergence times in great ape and human evolution. *Proc. Natl. Acad. Sci. U. S.*  
523 *A.* **109**, 15716–15721 (2012).
- 524 36. Meyer, M. *et al.* Nuclear DNA sequences from the Middle Pleistocene Sima de los  
525 Huesos hominins. *Nature* **531**, 504–507 (2016).
- 526 37. Posth, C. *et al.* Deeply divergent archaic mitochondrial genome provides lower time  
527 boundary for African gene flow into Neanderthals. *Nat. Commun.* **8**, (2017).
- 528 38. Gavryushkina, A. & Zhang, C. Total-Evidence Dating and the Fossilized Birth–Death  
529 Model. in *The Molecular Evolutionary Clock: Theory and Practice* (ed. Ho, S. Y. W.)  
530 175–193 (Springer International Publishing, 2020). doi:10.1007/978-3-030-60181-  
531 2\_11.
- 532 39. Will, M., Pablos, A. & Stock, J. T. Long-term patterns of body mass and stature  
533 evolution within the hominin lineage. *R. Soc. Open Sci.* **4**, (2017).
- 534 40. Grabowski, M. & Jungers, W. L. Evidence of a chimpanzee-sized ancestor of humans  
535 but a gibbon-sized ancestor of apes. *Nat. Commun.* **8**, 1–10 (2017).
- 536 41. Grabowski, M., Hatala, K. G. & Jungers, W. L. Body mass estimates of the earliest  
537 possible hominins and implications for the last common ancestor. *J. Hum. Evol.* **122**,  
538 84–92 (2018).



- 539 42. Grabowski, M. Bigger brains led to bigger bodies?: The correlated evolution of human  
540 brain and body size. *Curr. Anthropol.* **57**, 174–196 (2016).
- 541 43. Pagel, M. Modelling the evolution of continuously varying characters on phylogenetic  
542 trees: The case of Hominid cranial capacity. in *Morphology, Shape and Phylogeny*  
543 (eds. MacLeod, N. & Forey, P. L.) 269–286 (Taylor and Francis, London, 2002).  
544 doi:10.1201/9780203165171.
- 545 44. Gómez-Robles, A., Smaers, J. B., Holloway, R. L., Polly, P. D. & Wood, B. A. Brain  
546 enlargement and dental reduction were not linked in hominin evolution. *Proc. Natl.*  
547 *Acad. Sci. U. S. A.* **114**, 468–473 (2017).
- 548 45. Diniz-Filho, J. A. F., Jardim, L., Mondanaro, A. & Raia, P. Multiple Components of  
549 Phylogenetic Non-stationarity in the Evolution of Brain Size in Fossil Hominins. *Evol.*  
550 *Biol.* **46**, 47–59 (2019).
- 551 46. Miller, I. F., Barton, R. A. & Nunn, C. L. Quantitative uniqueness of human brain  
552 evolution revealed through phylogenetic comparative analysis. *Elife* **8**, (2019).
- 553 47. McHenry, H. M. & Coffing, K. *Australopithecus* to *Homo*: Transformations in Body  
554 and Mind. *Annu. Rev. Anthropol.* **29**, 125–146 (2000).
- 555 48. Grabowski, M., Voje, K. L. & Hansen, T. F. Evolutionary modeling and correcting for  
556 observation error support a 3/5 brain-body allometry for primates. *J. Hum. Evol.* **94**,  
557 106–116 (2016).
- 558 49. Montgomery, S. H., Capellini, I., Barton, R. A. & Mundy, N. I. Reconstructing the ups  
559 and downs of primate brain evolution: Implications for adaptive hypotheses and *Homo*  
560 *floresiensis*. *BMC Biol.* **8**, (2010).

- 561 50. Castiglione, S. *et al.* Ancestral State Estimation with Phylogenetic Ridge Regression.  
562 *Evol. Biol.* **47**, 220–232 (2020).
- 563 51. Potts, R. Environmental hypotheses of hominin evolution. *Am. J. Phys. Anthropol.*  
564 **107**, 93–136 (1998).
- 565 52. Potts, R. Evolution and climate variability. *Science* **273**, 922–923 (1996).
- 566 53. Zachos, J., Pagani, H., Sloan, L., Thomas, E. & Billups, K. Trends, rhythms, and  
567 aberrations in global climate 65 Ma to present. *Science* **292**, 686–693 (2001).
- 568 54. Richmond, B. G. & Jungers, W. L. *Orrorin tugenensis* femoral morphology and the  
569 evolution of hominin bipedalism. *Science* **319**, 1662–1665 (2008).
- 570 55. Crompton, R. H., Vereecke, E. E. & Thorpe, S. K. S. Locomotion and posture from the  
571 common hominoid ancestor to fully modern hominins, with special reference to the  
572 last common panin/hominin ancestor. *J. Anat.* **212**, 501–543 (2008).
- 573 56. Almécija, S. *et al.* The femur of *Orrorin tugenensis* exhibits morphometric affinities  
574 with both Miocene apes and later hominins. *Nat. Commun.* **4**, 1–12 (2013).
- 575 57. Kuperavage, A., Pokrajac, D., Chavanaves, S. & Eckhardt, R. B. Earliest known  
576 hominin calcar femorale in *Orrorin tugenensis* provides further Internal anatomical  
577 evidence for origin of human bipedal locomotion. *Anat. Rec.* **301**, 1834–1839 (2018).
- 578 58. Villmoare, B. *et al.* Early *Homo* at 2.8 Ma from Ledi-Geraru, Afar, Ethiopia. *Science*  
579 **347**, 1352–1355 (2015).
- 580 59. Harmand, S. *et al.* 3.3-million-year-old stone tools from Lomekwi 3, West Turkana,  
581 Kenya. *Nature* **521**, 310–315 (2015).

- 582 60. Leakey, M. G. *et al.* New hominin genus from eastern Africa shows diverse middle  
583 Pliocene lineages. *Nature* **410**, 433–440 (2001).
- 584 61. Brown, B., Brown, F. H. & Walker, A. New hominids from the Lake Turkana Basin,  
585 Kenya. *J. Hum. Evol.* **41**, 29–44 (2001).
- 586 62. Braun, D. R. *et al.* Earliest known Oldowan artifacts at >2.58 Ma from Ledi-Geraru,  
587 Ethiopia, highlight early technological diversity. *Proc. Natl. Acad. Sci. U. S. A.* **116**,  
588 11712–11717 (2019).
- 589 63. Lepre, C. J. *et al.* An earlier origin for the Acheulian. *Nature* **477**, 82–85 (2011).
- 590 64. Gamble, C. S. *Settling the Earth: the Archaeology of Deep Human History*.  
591 (Cambridge University Press: New York, 2013).
- 592 65. Hlubik, S., Berna, F., Feibel, C., Braun, D. & Harris, J. W. K. Researching the nature  
593 of fire at 1.5 Mya on the site of FxJj20 AB, Koobi Fora, Kenya, using high-resolution  
594 spatial analysis and FTIR spectrometry. *Curr. Anthropol.* **58**, S243–S257 (2017).
- 595 66. Hlubik, S. *et al.* Hominin fire use in the Okote member at Koobi Fora, Kenya: New  
596 evidence for the old debate. *J. Hum. Evol.* **133**, 214–229 (2019).
- 597 67. Gowlett, J. A. J. The discovery of fire by humans: A long and convoluted process.  
598 *Philos. Trans. R. Soc. B Biol. Sci.* **371**, 1–12 (2016).
- 599 68. Wrangham, Jones, Laden, Pilbeam & Conklin-Brittain. The Raw and the Stolen:  
600 Cooking and the Ecology of Human Origins. *Curr. Anthropol.* **40**, 567–594 (1999).
- 601 69. Fonseca-Azevedo, K. & Herculano-Houzel, S. Metabolic constraint imposes tradeoff  
602 between body size and number of brain neurons in human evolution. *Proc. Natl. Acad.*

- 603        *Sci. U. S. A.* **109**, 18571–18576 (2012).
- 604    70.    Carmody, R. N., Weintraub, G. S. & Wrangham, R. W. Energetic consequences of  
605        thermal and nonthermal food processing. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 19199–  
606        19203 (2011).
- 607    71.    Joordens, J. C. A. *et al.* *Homo erectus* at Trinil on Java used shells for tool production  
608        and engraving. *Nature* **518**, 228–231 (2015).
- 609    72.    Grün, R. *et al.* Dating the skull from Broken Hill, Zambia, and its position in human  
610        evolution. *Nature* **580**, 372–375 (2020).
- 611    73.    Timmermann, A. Quantifying the potential causes of Neanderthal extinction: Abrupt  
612        climate change versus competition and interbreeding. *Quat. Sci. Rev.* **238**, 106331  
613        (2020).
- 614    74.    Banks, W. E. *et al.* Neanderthal extinction by competitive exclusion. *PLoS One* **3**,  
615        e3972 (2008).
- 616    75.    Kochiyama, T. *et al.* Reconstructing the Neanderthal brain using computational  
617        anatomy. *Sci. Rep.* **8**, 1–9 (2018).
- 618    76.    Hershkovitz, I. *et al.* The earliest modern humans outside Africa. *Science* **359**, 456–  
619        459 (2018).
- 620    77.    Harvati, K. *et al.* Apidima Cave fossils provide earliest evidence of *Homo sapiens* in  
621        Eurasia. *Nature* **571**, 500–504 (2019).
- 622    78.    Hovers, E. Neandertals and modern humans in the middle Palaeolithic of the Levant:  
623        what kind of interaction? in *When Neandertals and Moderns Meet* (ed. Conard, N.)

- 624 65–85 (Kerns Verlag, Tübingen, 2006).
- 625 79. Scerri, E. M. L., Chikhi, L. & Thomas, M. G. Beyond multiregional and simple out-of-  
626 Africa models of human evolution. *Nat. Ecol. Evol.* **3**, 1370–1372 (2019).
- 627 80. Proctor, C., Douka, K., Proctor, J. W. & Higham, T. The Age and Context of the KC4  
628 Maxilla, Kent’s Cavern, UK. *Eur. J. Archaeol.* **20**, 74–97 (2017).
- 629 81. Villa, P. & Roebroeks, W. Neandertal demise: An archaeological analysis of the  
630 modern human superiority complex. *PLoS One* **9**, 96424 (2014).
- 631 82. Galway-Witham, J., Cole, J. & Stringer, C. Aspects of human physical and  
632 behavioural evolution during the last 1 million years. *J. Quat. Sci.* **34**, 355–378 (2019).
- 633 83. Downey, G. & Lende, D. H. Evolution and the brain. in *The Encultured Brain: an*  
634 *Introduction to Neuroanthropology* (eds. Lende, D. H. & Downey, G.) 103–138 (MIT  
635 Press: Cambridge, 2012).
- 636 84. Boivin, N. L. *et al.* Ecological consequences of human niche construction: Examining  
637 long-term anthropogenic shaping of global species distributions. *Proc. Natl. Acad. Sci.*  
638 *U. S. A.* **113**, 6388–6396 (2016).
- 639 85. Kendal, J., Tehrani, J. J. & Odling-Smee, J. Human niche construction in  
640 interdisciplinary focus. *Philos. Trans. R. Soc. B Biol. Sci.* **366**, 785–792 (2011).
- 641 86. Potts, R. Evolution and Environmental Change in Early Human Prehistory. *Annu. Rev.*  
642 *Anthropol.* **41**, 151–167 (2012).
- 643 87. Scerri, E. M. L. *et al.* Did Our Species Evolve in Subdivided Populations across  
644 Africa, and Why Does It Matter? *Trends Ecol. Evol.* **33**, 582–594 (2018).

- 645 88. Sol, D., Bacher, S., Reader, S. M. & Lefebvre, L. Brain size predicts the success of  
646 mammal species introduced into novel environments. *Am. Nat.* **172**, 63–71 (2008).
- 647 89. Potts, R. Hominin evolution in settings of strong environmental variability. *Quat. Sci.*  
648 *Rev.* **73**, 1–13 (2013).
- 649 90. Wedage, O. *et al.* Specialized rainforest hunting by *Homo sapiens* ~45,000 years ago.  
650 *Nat. Commun.* **10**, 1–8 (2019).
- 651 91. Roberts, P. & Stewart, B. A. Defining the ‘generalist specialist’ niche for Pleistocene  
652 *Homo sapiens*. *Nat. Hum. Behav.* **2**, 542–550 (2018).
- 653 92. Van Heteren, A. H. & Sankhyan, A. R. Hobbits and pygmies: trends in evolution. in  
654 *Asian perspectives on human evolution* (ed. Sankhyan, A. R.) 172–187 (Serials  
655 Publications, New Delhi, India, 2009).
- 656 93. Diniz-Filho, J. A. F. & Raia, P. Island rule, quantitative genetics and brain-body size  
657 evolution in *Homo floresiensis*. *Proc. R. Soc. B Biol. Sci.* **284**, 20171065 (2017).
- 658 94. Kubo, D., Kono, R. T. & Kaifu, Y. Brain size of *Homo floresiensis* and its  
659 evolutionary implications. *Proc. R. Soc. B Biol. Sci.* **280**, (2013).
- 660 95. Dennell, R. W., Louys, J., O’Regan, H. J. & Wilkinson, D. M. The origins and  
661 persistence of *Homo floresiensis* on Flores: Biogeographical and ecological  
662 perspectives. *Quat. Sci. Rev.* **96**, 98–107 (2014).
- 663 96. Foster, J. B. Evolution of mammals on Islands. *Nature* **202**, 234–235 (1964).
- 664 97. Brown, P. *et al.* A new small-bodied hominin from the Late Pleistocene of Flores,  
665 Indonesia. *Nature* **431**, 1055–1061 (2004).

- 666 98. Argue, D., Groves, C. P., Lee, M. S. Y. & Jungers, W. L. The affinities of *Homo*  
667 *floresiensis* based on phylogenetic analyses of cranial, dental, and postcranial  
668 characters. *J. Hum. Evol.* **107**, 107–133 (2017).
- 669 99. Brumm, A. *et al.* Early stone technology on Flores and its implications for *Homo*  
670 *floresiensis*. *Nature* **441**, 624–628 (2006).
- 671 100. Moore, M. W. & Brumm, A. *Homo floresiensis* and the African Oldowan. in  
672 *Interdisciplinary Approaches to the Oldowan* (eds. Hovers, E. & Braun, D. R.) 61–69  
673 (Springer, New York, 2009). doi:10.1007/978-1-4020-9060-8\_6.
- 674 101. Garvin, H. M. *et al.* Body size, brain size, and sexual dimorphism in *Homo naledi*  
675 from the Dinaledi Chamber. *J. Hum. Evol.* **111**, 119–138 (2017).
- 676 102. Laird, M. F. *et al.* The skull of *Homo naledi*. *J. Hum. Evol.* **104**, 100–123 (2017).
- 677 103. Holloway, R. L. *et al.* Endocast morphology of *Homo naledi* from the Dinaledi  
678 Chamber, South Africa. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 5738–5743 (2018).
- 679 104. Kivell, T. L. *et al.* The hand of *Homo naledi*. *Nat. Commun.* **6**, 1–9 (2015).
- 680 105. Marchi, D. *et al.* The thigh and leg of *Homo naledi*. *J. Hum. Evol.* **104**, 174–204  
681 (2017).
- 682 106. Dirks, P. H. *et al.* The age of *Homo naledi* and associated sediments in the Rising Star  
683 Cave, South Africa. *Elife* **6**, e24231 (2017).
- 684 107. Aiello, L. C. & Wheeler, P. The Expensive-Tissue Hypothesis: The Brain and the  
685 Digestive System in Human and Primate Evolution. *Curr. Anthropol.* **36**, 199–221  
686 (1995).

- 687 108. Isler, K. & van Schaik, C. P. The Expensive Brain: A framework for explaining  
688 evolutionary changes in brain size. *J. Hum. Evol.* **57**, 392–400 (2009).
- 689 109. Isler, K. & Van Schaik, C. P. Metabolic costs of brain size evolution. *Biol. Lett* **2**, 557–  
690 560 (2006).
- 691 110. Berthaume, M. A., Delezenne, L. K. & Kupczik, K. Dental topography and the diet of  
692 *Homo naledi*. *J. Hum. Evol.* **118**, 14–26 (2018).
- 693 111. Nascimento, F. F., Reis, M. Dos & Yang, Z. A biologist’s guide to Bayesian  
694 phylogenetic analysis. *Nat. Ecol. Evol.* **1**, 1446–1454 (2017).
- 695 112. O’Reilly, J. E., dos Reis, M. & Donoghue, P. C. J. Dating Tips for Divergence-Time  
696 Estimation. *Trends Genet.* **31**, 637–650 (2015).
- 697 113. Yang, Z. & Donoghue, P. C. J. Dating species divergences using rocks and clocks.  
698 *Philos. Trans. R. Soc. B Biol. Sci.* **371**, 20150126 (2016).
- 699 114. Cameron, D. W. & Groves, C. P. *Bones, Stones and Molecules: Out of Africa and*  
700 *Human Origins*. (Elsevier, San Diego, 2004).
- 701 115. Cameron, D., Patnaik, R. & Sahni, A. The phylogenetic significance of the middle  
702 pleistocene Narmada hominin cranium from central India. *Int. J. Osteoarchaeol.* **14**,  
703 419–447 (2004).
- 704 116. Berger, L. R. *et al.* *Australopithecus sediba*: A New Species of Homo-Like  
705 Australopith from South Africa. *Science* **328**, 195–204 (2010).
- 706 117. Irish, J. D., Guatelli-Steinberg, D., Legge, S. S., De Ruiter, D. J. & Berger, L. R.  
707 Dental morphology and the phylogenetic ‘place’ of *Australopithecus sediba*. *Science*



- 708           **340**, 1233062 (2013).
- 709   118.   Lordkipanidze, D. *et al.* A complete skull from Dmanisi, Georgia, and the evolutionary  
710           biology of early *Homo*. *Science* **342**, 326–331 (2013).
- 711   119.   Kimbel, W. H., Rak, Y. & Johanson, D. C. *The skull of Australopithecus afarensis*.  
712           (Oxford University Press, New York, 2004). doi:10.5860/choice.42-1028.
- 713   120.   Strait, D. S. & Grine, F. E. Inferring hominoid and early hominid phylogeny using  
714           craniodental characters: The role of fossil taxa. *J. Hum. Evol.* **47**, 399–452 (2004).
- 715   121.   Chang, M. L. Neandertal origins, Middle Pleistocene systematics, and tests of current  
716           taxonomic and phylogenetic hypotheses. Ph.D. Dissertation, University of  
717           Pennsylvania. (2005).
- 718   122.   Martínón-Torres, M. *et al.* Dental evidence on the hominin dispersals during the  
719           Pleistocene. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 13279–13282 (2007).
- 720   123.   Gilbert, W. H. *Homo erectus* cranial anatomy. in *Homo erectus: Pleistocene Evidence*  
721           *from the Middle Awash, Ethiopia* (eds. Gilbert, W. H. & Asfaw, B.) 265–311  
722           (University of California Press, Berkeley, 2008).
- 723   124.   Argue, D., Morwood, M. J., Sutikna, T., Jatmiko & Saptomo, E. W. *Homo floresiensis*:  
724           a cladistic analysis. *J. Hum. Evol.* **57**, 623–639 (2009).
- 725   125.   Mounier, A., Marchal, F. & Condemi, S. Is *Homo heidelbergensis* is a distinct species?  
726           New insight on the Mauer mandible. *J. Hum. Evol.* **56**, 219–246 (2009).
- 727   126.   Zeitoun, V. *The Human Canopy: Homo erectus, Homo soloensis, Homo pekinensis*  
728           *and Homo floresiensis*. (B.A.R. S1937, 2009).

- 729 127. Mongle, C. S., Strait, D. S. & Grine, F. E. Expanded character sampling underscores  
730 phylogenetic stability of *Ardipithecus ramidus* as a basal hominin. *J. Hum. Evol.* **131**,  
731 28–39 (2019).
- 732 128. Martin, J. M. *et al.* Drimolen cranium DNH 155 documents microevolution in an early  
733 hominin species. *Nat. Ecol. Evol.* **5**, 38–45 (2021).
- 734 129. Finstermeier, K. *et al.* A Mitogenomic Phylogeny of Living Primates. *PLoS One* **8**,  
735 69504 (2013).
- 736 130. Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular  
737 evolutionary genetics analysis across computing platforms. *Mol. Biol. Evol.* **35**, 1547–  
738 1549 (2018).
- 739 131. Lanfear, R., Frandsen, P. B., Wright, A. M., Senfeld, T. & Calcott, B. Partitionfinder  
740 2: New methods for selecting partitioned models of evolution for molecular and  
741 morphological phylogenetic analyses. *Mol. Biol. Evol.* **34**, 772–773 (2017).
- 742 132. Lanfear, R., Calcott, B., Ho, S. Y. W. & Guindon, S. PartitionFinder: Combined  
743 selection of partitioning schemes and substitution models for phylogenetic analyses.  
744 *Mol. Biol. Evol.* **29**, 1695–1701 (2012).
- 745 133. Lewis, P. A Likelihood Approach to Estimating Phylogeny from Discrete  
746 Morphological Character Data. *Soc. Syst. Biol.* **50**, 913–925 (2001).
- 747 134. Püschel, H. P., O'Reilly, J. E., Pisani, D. & Donoghue, P. C. J. The impact of fossil  
748 stratigraphic ranges on tip-calibration, and the accuracy and precision of divergence  
749 time estimates. *Palaeontology* **63**, 67–83 (2020).
- 750 135. Katoh, S. *et al.* New geological and palaeontological age constraint for the gorilla-

751 human lineage split. *Nature* **530**, 215–218 (2016).

752 136. Kanimatsu, Y. *et al.* A new Late Miocene great ape from Kenya and its implications  
753 for the origins of African great apes and humans. *Proc. Natl. Acad. Sci. U. S. A.* **104**,  
754 19220–19225 (2007).

755 137. Begun, D. R. Miocene Hominids and the Origins of the African Apes and Humans.  
756 *Annu. Rev. Anthropol.* **39**, 67–84 (2010).

757 138. Begun, D. R., Nargolwalla, M. C. & Kordos, L. European Miocene Hominids and the  
758 Origin of the African Ape and Human Clade. *Evol. Anthropol.* **21**, 10–23 (2012).

759 139. Suwa, G., Kono, R. T., Katoh, S., Asfaw, B. & Beyene, Y. A new species of great ape  
760 from the late Miocene epoch in Ethiopia. *Nature* **448**, 921–924 (2007).

761 140. Harrison, T. Dendropithecoidae, Proconsuloidae, and Hominoidea. in *Cenozoic*  
762 *mammals of Africa* (eds. Werdelin, L. & Sanders, W.) 429–470 (University of  
763 California Press, Berkeley, 2012). doi:10.1525/california/9780520257214.003.0024.

764 141. Fu, Q. *et al.* Genome sequence of a 45,000-year-old modern human from western  
765 Siberia. *Nature* **514**, 445–449 (2014).

766 142. Fu, Q. *et al.* A revised timescale for human evolution based on ancient mitochondrial  
767 genomes. *Curr. Biol.* **23**, 553–559 (2013).

768 143. Berger, L. R., Hawks, J., Dirks, P. H. G. M., Elliott, M. & Roberts, E. M. *Homo naledi*  
769 and Pleistocene hominin evolution in subequatorial Africa. *Elife* **6**, e24234 (2017).

770 144. Kimbel, W. H. & Rak, Y. *Australopithecus sediba* and the emergence of *Homo*:  
771 Questionable evidence from the cranium of the juvenile holotype MH 1. *J. Hum. Evol.*

772           **107**, 94–106 (2017).

773   145.   Schroeder, L. *et al.* Skull diversity in the *Homo* lineage and the relative position of  
774           *Homo naledi*. *J. Hum. Evol.* **104**, 124–135 (2017).

775   146.   Ronquist, F. *et al.* Mrbayes 3.2: Efficient bayesian phylogenetic inference and model  
776           choice across a large model space. *Syst. Biol.* **61**, 539–542 (2012).

777   147.   Miller, M. A., Pfeiffer, W. & Schwartz, T. Creating the CIPRES Science Gateway for  
778           inference of large phylogenetic trees. in *Proceedings of the Gateway Computing*  
779           *Enviroments Workshop (GCE)* 1–8 (New Orleans,LA, 2010).  
780           doi:10.1109/GCE.2010.5676129.

781   148.   Rambaut, A., Drummond, A. J., Xie, D., Baele, G. & Suchard, M. A. Posterior  
782           Summarization in Bayesian Phylogenetics Using Tracer 1.7. *Syst. Biol.* **67**, 901–904  
783           (2018).

784   149.   Zhang, C., Stadler, T., Klopstein, S., Heath, T. A. & Ronquist, F. Total-Evidence  
785           Dating under the Fossilized Birth-Death Process. *Syst. Biol.* **65**, 228–249 (2016).

786   150.   Jerison, H. J. *Evolution of the Brain and Intelligence*. (Academic Press, New York,  
787           1973).

788   151.   Jerison, H. J. Brain, body and encephalization in early primates. *J. Hum. Evol.* **8**, 615–  
789           635 (1979).

790   152.   Cairó, O. External measures of cognition. *Front. Hum. Neurosci.* **5**, 108 (2011).

791   153.   Stephan, H., Frahm, H. D. & Baron, G. New and revised data on volumes of brain  
792           structures in insectivores and primates. *Folia Primatol.* **35**, 1–29 (1981).

154. R Core Team. *R: A language and environment for statistical computing*. (R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>, 2019).
155. Paradis, E., Claude, J. & Strimmer, K. APE: Analyses of phylogenetics and evolution in R language. *Bioinformatics* **20**, 289–290 (2004).
156. Pinheiro, J., Bates, D., DebRoy, S. & Sarkar, D. nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-147. (2020).
157. Ives, A. R. R 2 s for Correlated Data: Phylogenetic Models, LMMs, and GLMMs. *Syst. Biol.* **68**, 234–251 (2019).
158. Revell, L. J. phytools: An R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* **3**, 217–223 (2012).
159. Püschel, H. P., Bertrand, O. C., O'Reilly, J. E., Bobe, R. & Püschel, T. A. Divergence-time estimates for hominins provide insight into encephalization and body mass trends in human evolution: Data and Code. GitHub.  
<https://github.com/HansPueschel/Hominin-div-time-evolution>;  
<https://dx.doi.org/10.5281/zenodo.4537445>. (2021).

## Acknowledgements

This research was funded by the National Agency for Research and Development (ANID)/PFCHA/Doctorado en el extranjero Becas Chile/2018-72190003 to H.P.P; a Marie Skłodowska-Curie Actions: Individual Fellowship (H2020-MSCA-IF-2018-2020; No. 792611) and the European Research Council (ERC) Starting Grant PalM, under the European

814 Union's Horizon 2020 Research and Innovation Programme (No. 756226) to O.C.B.; the  
815 Leverhulme Trust Early Career Fellowship, ECF-2018-264 to T.A.P.

## 816 **Author contributions**

817 H.P.P. and T.A.P. conceived and designed the study. O.C.B. compiled the body mass and  
818 endocranial volume dataset. J.E.O. provided methodological support. H.P.P. and T.A.P.  
819 carried out all the mentioned analyses and wrote an initial draft. H.P.P., O.C.B., J.E.O., R.B.  
820 and T.A.P. interpreted the obtained results and contributed to the writing of the submitted  
821 version of this work.

## 822 **Competing interests**

823 The authors declare no competing interests.

## 824 **Figures**

825 **Figure 1.** Alternative topological hypotheses tested in the total-evidence dating (TED)  
826 analyses. a), similar to the phylogeny of Dembo *et al.*<sup>17</sup>; b), the same topology as a) but  
827 moving *Au. sediba* from the *Homo* clade, to be the sister taxa to *Au. africanus*; c), the same  
828 topology as a) but changing the position of *H. naledi* from the stem of *H. antecessor* to be the  
829 sister taxa to the African *H. erectus*; d), the same topology of a) but taking *H. floresiensis*  
830 from the base of the genus *Homo* to be the sister taxa to the Asian *H. erectus*.

831 **Figure 2.** Summary diagram of important paleoclimatic and hominin evolution events plotted  
832 next to the four obtained consensus phylogenies and time divergence estimates (Red node  
833 bars represent the 95% highest posterior density [HPD] for the estimated node ages). a)  
834 Dembo *et al.*<sup>17</sup> hypothesis, b) *Au. sediba* hypothesis, c) *H. naledi* hypothesis, and d) *H.*  
835 *floresiensis* hypothesis. A composite benthic foraminifera oxygen isotope record obtained  
836 from<sup>53</sup> is displayed at the top of the figure to illustrate the evolution of high-latitude glacial

cycles and dominant periodicity of glacial variability, as well as palaeotemperatures (the red line corresponds to a smoothing spline used to depict the main trend in  $\delta^{18}\text{O}$  values; smoothing parameter=0.2).

**Figure 3.** Body mass ancestral character state reconstructions (ACSR) for each species mapped onto the four consensus time-calibrated phylogenies. a) Dembo *et al.*<sup>17</sup> hypothesis, b) *Au. sediba* hypothesis, c) *H. naledi* hypothesis, and d) *H. floresiensis* hypothesis. The values at nodes and branches were reconstructed using a maximum-likelihood ancestral character estimation method under a Brownian motion model.

**Figure 4.** PEQ ancestral character state reconstructions (ACSR) for each species mapped onto the four consensus time-calibrated phylogenies. a) Dembo *et al.*<sup>17</sup> hypothesis, b) *Au. sediba* hypothesis, c) *H. naledi* hypothesis, and d) *H. floresiensis* hypothesis. The values at nodes and branches were reconstructed using a maximum-likelihood ancestral character estimation method under a Brownian motion model.

**Figure 5.** Boxplots of a-d) body mass (kg) and e-h) PEQ ancestral character state reconstructions (ACSR) per node based on a sample of 9002 time-calibrated posterior trees for each of the tested hypothesis. a, e) Dembo *et al.*<sup>17</sup> hypothesis, b, f) *Au. sediba* hypothesis, c, g) *H. naledi* hypothesis, and d, h) *H. floresiensis* hypothesis. The red dots indicate the ACSR conducted using the consensus trees. The median is indicated by the horizontal black line, the interquartile range (IQR) is the white box, and the whiskers indicate the minimum and the maximum (at 1.5 \* IQR of the lower and upper hinge respectively).

**Table 1.** Divergence times mean and 95% highest posterior density interval (HPD) in Ma for the different phylogenetic hypotheses in Fig. 1. The maximum and minimum bounds for the 95% HPD are in parentheses. Abbreviations: Div., divergence-times mean; d, Dembo *et al.*<sup>17</sup> hypothesis; s, *Au. sediba* hypothesis; n, *H. naledi* hypothesis; f, *H. floresiensis* hypothesis.

Node*	Div. mean-d	Div. mean-s	Div. mean-n	Div. mean-f
1	10.35 (11.40, 10.00)	10.35 (11.42, 10.00)	10.37 (11.49, 10.00)	10.33 (11.36, 10.00)
2	7.47 (8.52, 6.66)	7.43 (8.43, 6.61)	7.50 (8.59, 6.67)	7.46 (8.53, 6.63)
3	7.20 (8.11, 6.51)	7.15 (8.06, 6.49)	7.22 (8.15, 6.51)	7.18 (8.10, 6.47)
4	6.40 (7.59, 5.21)	6.41 (7.55, 5.26)	6.42 (7.57, 5.24)	6.39 (7.54, 5.21)
5	5.44 (6.49, 4.60)	5.44 (6.42, 4.61)	5.50 (6.52, 4.64)	5.42 (6.47, 4.56)
6	4.90 (5.76, 4.18)	4.97 (5.80, 4.24)	4.95 (5.82, 4.24)	4.87 (5.72, 4.15)
7	3.97 (4.47, 3.63)	3.94 (4.40, 3.62)	3.98 (4.50, 3.63)	3.97 (4.46, 3.62)
8	4.52 (5.32, 3.83)	4.60 (5.41, 3.92)	4.56 (5.37, 3.86)	4.49 (5.29, 3.82)
9	4.06 (4.76, 3.44)	4.12 (4.78, 3.52)	4.09 (4.80, 3.46)	4.02 (4.73, 3.39)
10	3.83 (4.51, 3.19)	3.93 (4.58, 3.34)	3.85 (4.57, 3.19)	3.77 (4.46, 3.13)
11	2.93 (3.34, 2.65)	2.93 (3.33, 2.65)	2.94 (3.36, 2.66)	2.93 (3.34, 2.65)
12	2.47 (2.81, 2.22)	2.46 (2.79, 2.22)	2.48 (2.82, 2.22)	2.47 (2.81, 2.22)
13	3.47 (4.21, 2.75)	3.24 (4.05, 2.59)	3.52 (4.30, 2.81)	<b>1.73 (2.09, 1.47)</b>
14	2.94 (3.46, 2.53)	2.81 (3.30, 2.40)	2.98 (3.50, 2.55)	3.04 (3.62, 2.56)
15	2.70 (3.17, 2.35)	<b>3.57 (4.23, 2.94)</b>	2.73 (3.21, 2.36)	2.75 (3.25, 2.34)
16	2.63 (3.10, 2.25)	2.54 (2.97, 2.19)	2.67 (3.15, 2.28)	2.70 (3.21, 2.26)
17	2.39 (2.83, 2.02)	2.33 (2.73, 1.99)	2.44 (2.88, 2.06)	2.45 (2.93, 2.05)
18	2.13 (2.54, 1.79)	2.09 (2.47, 1.78)	2.18 (2.58, 1.82)	2.19 (2.63, 1.83)
19	1.89 (2.27, 1.57)	1.87 (2.21, 1.57)	1.91 (2.30, 1.57)	1.98 (2.37, 1.64)
20	1.59 (2.02, 1.20)	1.57 (1.99, 1.20)	<b>1.94 (2.35, 1.58)</b>	1.61 (2.06, 1.18)
21	1.42 (1.83, 1.04)	1.42 (1.81, 1.04)	1.62 (2.09, 1.17)	1.43 (1.84, 1.01)
22	1.09 (1.50, 0.73)	1.08 (1.46, 0.74)	1.17 (1.59, 0.76)	1.08 (1.50, 0.70)
23	0.77 (1.13, 0.40)	0.77 (1.12, 0.42)	0.80 (1.20, 0.43)	0.76 (1.16, 0.38)
24	0.60 (1.01, 0.26)	0.59 (0.94, 0.27)	0.63 (1.03, 0.28)	0.59 (1.02, 0.25)

\*Node's ages that are not comparable due to the specific changes in the phylogenetic hypotheses are indicated in bold.